

CATEGORIES, CONTROLS AND CEILINGS LAWS REPLY

Keith R. Laws

(Brain and Cognition Research Group, Division of Psychology, Nottingham Trent University, UK)

In the target paper, I outlined several methodological issues associated with attempts to document category specific deficits; and a potential solution based around certain minimal criteria. The main argument being that an accurate interpretation of patient performance requires a comparison with a normal control group and that the group performs below ceiling. Neither of these requirements is new or one would imagine, especially contentious. It is therefore surprising that a review of the category specific literature reveals no single study that meets these criteria (Laws, in press). Moreover, this has to be viewed in the context that the commonly used analyses (within-patient χ^2 and between-subject comparisons with controls at ceiling) produce false positive, false negative and even *paradoxical* deficits.

The target paper commentaries may be roughly classified into those on the one hand in favour of using controls who are not at ceiling (Sartori and Lombardi; McMullen and Filliter; Rosazza et al.) and that analyses of healthy controls are critical in themselves (Låg; Marques), through to those who feel that their data already meets the criteria in the target paper (Capitani and Laiacona; Laiacona and Barbarotto; Bright et al.) or argue that controls are not necessarily required (Marshall and Gurd's extreme cases approach). This reply will focus primarily on general points of contention raised by the contributors.

Solutions to the Ceiling Effect in Controls

Capitani and Laiacona discuss various possible solutions to address ceiling effects in controls, including: degrading the stimuli in some way to make them more difficult for controls, using controls who perform below ceiling (for example, older subjects)ⁱ or constructing a more difficult stimulus set.

Both Capitani and Laiacona and Marshall and Gurd reject the stimulus degradation idea because it would introduce unknown variables (see Låg) as also remarked upon in the target article (see comments on Turnbull and Laws, 2000). Although

stimulus degradation is a useful tool for examining the factors affecting the category differences in healthy subjects (Låg; Marques; Låg, in press; Laws and Neve 1999; Laws et al., 2002; Laws, 2000), it cannot provide control baselines. This becomes apparent when one considers that different presentation conditions may interact with category (see Låg, in press; Gerlach, 2001; Laws and Neve, 1999).

The solution preferred by Capitani and Laiacona is a "slightly different" group of normal subjects who in this case, consist of 60 elderly subjects (30 males and 30 females aged 71). These elderly subjects used in all of the studies by Capitani, Laiacona and colleagues do however have naming at ceiling with $91.83 \pm 12.8\%$ and $96.38 \pm 7.33\%$ of the controls naming all of the living things and nonliving things correctlyⁱⁱ.

Like Capitani and Laiacona, Marshall and Gurd have concerns about using 'more difficult' stimuli because of possible 'floor effects' in patients and because the stimuli might be "more subject to any peculiar habits of the subjects". Graded naming tests, however, need not necessarily engender floor effects; and for example, Laws et al. (in press) reported picture naming for Alzheimer patients at 25% (not at floor), while controls performed at 75% (not at ceiling). Similarly, Rosazza et al., document a living deficit in their patient using a variation on this approach. Concerning the point about "peculiar habits", it is of course, inevitable that graded tests would more successfully tap and reflect individual variability. Indeed, part of the attempt to explain category effects should be directed toward understanding normal individual variability (rather than homogenous perfect performance).

Normal Individual Variability

As noted by Marques, natural asymmetries of category processing exist in healthy subjects and so, control data are essential to "...even establish

ⁱ Another unmentioned possibility might be to use pathological controls to attain a below ceiling 'control'. Such an approach may indicate abnormality, but critically not the direction. This is exemplified by the problems associated with double dissociations (see Figure 1) that are not referenced to healthy control data.

ⁱⁱ More critically, while the regression method advocated by Capitani and colleagues could potentially address some of the issues raised in the target paper, it is not immune to ceiling effects. In the original article, I mistook the term 'Name Agreement' used by Capitani, Laiacona and colleagues' papers to mean what it typically means, i.e. the number of alternative names produced rather than an indicator that control data were included in the regression.




Fig. 1 – Category double dissociation or two single dissociations of the same category?

Legend. The figure on the left shows a typical double dissociation between living and nonliving things; however, these are unreferenced to control data. The figure on the right shows an alternative possibility with fake control data added. Contrary to the implication from the double dissociation on the left, this shows that both patient 1 and 2 are impaired for Living things and are normal for Nonliving things. Of course, the control data could be drawn to provide the converse example.

that a real deficit is present". Therefore, as Sartori and Lombardi remark, we cannot rely upon generic control groups because category performance may be affected by sex, education and age differences. In the context of normal individual variability, Laiacona and Barbarotto refer to several studies documenting an interaction between sex and category (e.g. Capitani et al., 1999; Barbarotto et al., 2002; Laws, 1999, 2003) with males performing better with nonliving things and females with living thingsⁱⁱⁱ. Findings relating to such individual differences are important both as an aid to understanding category effects and in determining the nature of appropriate controls i.e. they indicate that controls should be sex matched.

Illusions of Double Dissociations beyond Naming

An altogether different issue raised by Laiacona and Barbarotto and by Bright et al. concerns the emphasis on picture naming in the target article. While picture naming does represent just one test of category specificity, it was chosen for a variety of reasons, including the fact that it is the one test common to almost all studies and is used (implicitly or otherwise) as the standard means to documenting category effects. Of course, most studies use many other tests; however, the choice of any other single test is likely to be less informative and less widely used (e.g. drawing, naming-to-description, sentence verification).

If one accepts the methodological problems relating to controls in picture naming studies, it would be unsurprising to see that the same problems extend to other tests in the same studies (or even beyond category specificity as alluded to by Marshall and Gurd). Indeed, a cursory glance at category specific studies (or the wider literature) appears to confirm this suspicion. As an example, let us examine the semantic question data for PL and MF (Laiacona and Barbarotto)^{iv} as a potential double dissociation across living and nonliving things. The absolute scores and the cross-over figure do look convincing, and are typical of the evidence presented for double dissociations (and ultimately the fractionation of cognition). Nevertheless, the normative data for this task come from the same 60 elderly individuals and is again at ceiling (96.7% and 98.6% of subjects answered every living and nonliving thing question correctly). Therefore, the ceiling effect problem with naming also extends to other tests in category specific investigations. At a more abstract level, consider the typical double dissociation in this area of enquiry (which is unreferenced to controls) below and how the interpretation might change when control data are added. This example shows how absolute performance may be misleading. An apparent and typical double dissociation, when referenced to controls, may reveal the same category deficit in both patients (here for living things).

ⁱⁱⁱ Marshall and Gurd provide an example in which it seems John Marshall knows nonliving things such as demijohn and tantalus, but not armadillo and platypus. One possible explanation is the interaction between category and sex.

^{iv} Capitani and colleagues refer to several of their cases that "escaped my attention" (re: Barbarotto et al., 1995 [MF]; Albanese et al., 2000 [GR and PL]; Laiacona et al., 2003 [EA]). Three of these cases were actually referenced in the target paper – each being reported on more than one occasion elsewhere: PL (Laiacona and Capitani, 2001); GR (Laiacona et al., 1993); EA (Barbarotto et al., 1996; Laiacona et al., 1997).

Extreme Cases

Marshall and Gurd refer to the method of 'extreme cases', where patients, but not normal subjects for example "...make semantic errors when reading individual words...(or) draw clockfaces that only include the numbers 12, 1, 2, 3, 4, 5 and 6". These examples, however, are dealing with *qualitatively* different human behaviours. While in category specificity (and many other neuropsychological deficits), the focus is on quantitative differences and it is not known *a priori* whether normal subjects would show a living or nonliving advantage. In this sense, while *extreme cases* may be useful, they are unusual and in the minority of published neuropsychological case studies.

If Normal Subjects show a Living Advantage, then why are there so many Living Deficits?

Bright et al. suggest that some of the points I raised in the target article are "ill-judged and contradictory". The focus of the argument by Bright and colleagues appears to rest on two assumptions that they take to be contradictory: (a) the finding that normals show better naming of living than nonliving things; and (b) the far greater incidence of living deficits reported in patients.

Turning to the first assumption, if materials are matched across category on the usual nuisance variables, controls do show better naming of living than nonliving things (see Laws, 1999, 2000, 2002a, 2003; Laws et al., 2002b; Laws and Neve, 1999)^v. Despite the assumptions that some researchers build into their models of category specificity, I am not aware of any study of normal subjects reporting a naming advantage for nonliving things on matched stimuli^{vi}. The second assumption is one that I have directly questioned (Laws et al., in press; Laws, in press). Indeed, Laws et al. (in press) have shown that over-reliance on stimuli that engender ceiling effects in controls may inflate the proportion of living cases and underestimate the number of nonliving ones (Laws et al., in press; Laws et al., 2003)^{vii}.

Bright et al. suggest that that I have been over-dismissive of their data for JBR (Bunn et al., 1998)

and RC (Moss et al., 1998). In both cases, the same control data were taken from a colour photo set in 40 young controls and 8 elderly controls. Contrary to what Bright et al. claim, both control groups were at ceiling (living and nonliving things, young controls score 93 vs. 96% correct; and for the 8 elderly controls score 95 vs. 95% correct).

Finally, they refer to the contradictory findings from the case of SE (originally described by Laws et al. (1995), but later also described by Moss et al., 1998; see also Laws, 1998). If a patient produces contradictory results across groups, then it does seem grounds for doubting the findings of both studies and a warning about comparing across different paradigms and materials (see also commentary by Låg; see also Laws, 1998). Second, and critically, they argue that they "...did *meet all of Laws' criteria* for comparison to control groups" (my italics), but this is not the case. Three sets of naming data are presented: the full Snodgrass and Vanderwart corpus (using 12 controls) of stimuli unmatched across category; a subset of those stimuli matched for frequency and familiarity, but with no controls; and the colour photos from Bunn et al.; here used a within patient comparison and no reference to controls. Hence, none of their data meets the criteria outlined in the target article.

Leaving aside Laws et al. (in press) and Rosazza et al., which demonstrate the approach advocated, and despite the claims made in some commentaries, no study has documented a category effect that meets the criteria outlined in the target article. In other words, none have demonstrated a category effect using: (a) a matched (sex, age etc) control group who; (b) do not perform at ceiling on; (c) stimuli that have the usual cross-category confounding differences controlled; and (d) in which the sample statistics are treated *as* statistics rather than as population parameters^{viii}. Moreover, this means that, currently, no soundly established double dissociation between living and nonliving things exists. As noted in the target paper, this does not mean that category effects do not exist. Indeed, the approach outlined above represents a set of minimal criteria that will enhance the accuracy of their documentation and interpretation. While it remains to be determined whether these control problems apply more generally in cognitive neuropsychology (pace Marshall and Gurd), they do extend beyond picture naming and pervasively affect category specific studies.

REFERENCES

- ALBANESE E, CAPITANI E, BARBAROTTO R and LAIACONA M. Semantic category dissociations, familiarity and gender. *Cortex*, 36: 733–746, 2000.

^v As an aside, this would also not fit with the assumption of McMullen and Filliter concerning the notion that because living things typically come from fewer categories than nonliving things, this creates a disadvantage for the former (this rests on the assumption of greater structural overlap – which has been doubted: see Laws and Gale, 2002). Indeed, it could also be equally well argued that by circumscribing the categories for living things, it provides an advantage for them.

^{vi} Nonetheless, it is, of course, possible that normal subjects will show an advantage for nonliving things in some circumstances (Låg, in press).

^{vii} Finally, there is no necessary contradiction in the premises that normal subjects name more living than nonliving things and that the ratio of living deficits is greater than for nonliving. Both could, for example, reflect the presence of a specific neural substrate dedicated to processing living things (while no single system exists for nonliving things). Hence, a dedicated system would aid normal subjects in processing living things easier and be more susceptible to damage (See Laws, 1999).

^{viii} See Crawford and Garthwaite (2002) for a detailed description of the statistical issues and Laws, Gale, Leeson and Crawford (in press) for examples of the use of these statistics to category deficits in Alzheimer's patients.

- BARBAROTTO R, LAIACONA M, MACCHI V and CAPITANI E. Picture reality decision, semantic categories and gender – a new set of pictures, with norms and an experimental study. *Neuropsychologia*, 40: 1637-1653, 2002.
- BARBAROTTO R, CAPITANI E and LAIACONA M. Naming deficit in herpes simplex encephalitis. *Acta Neurologica Scandinavica*, 93: 272-280, 1996.
- BRIGHT P, MOSS HE and TYLER LK. Commentary on Keith R Laws: "Illusions of Normality". *Cortex*, 41: 2005.
- BUNN EM, TYLER LK and MOSS HE. Category-specific semantic deficits: The role of familiarity and property type reexamined. *Neuropsychology*, 12: 367-379, 1998a.
- CAPITANI E and LAIACONA M. An illusory illusion? Comments on "Illusions of Normality" by Keith Laws. *Cortex*, 41: 2005.
- CAPITANI E, LAIACONA M and BARBAROTTO R. Gender affects word retrieval of certain categories in semantic fluency tasks. *Cortex*, 35: 273-278, 1999.
- CRAWFORD JR and GARTHWAITE PH. Investigation of the single case in neuropsychology: Confidence limits on the abnormality of test scores and test score differences. *Neuropsychologia*, 40: 1196-1208, 2002.
- GERLACH C. Structural similarity causes different category-effects depending on task characteristics. *Neuropsychologia*, 39: 895-900, 2001.
- LÅG T. Illusions of category specificity? A commentary on Keith Laws' forum article. *Cortex*, 41: 2005.
- LÅG T. (in press). Category-specific effects in object identification: What is 'normal'. *Cortex*, 41: 2005.
- LAIACONA M and BARBAROTTO R. On double dissociations, controls and gender: Some neglected data about category specificity. *Cortex*, 41: 2005.
- LAIACONA M, BARBAROTTO R and CAPITANI E. Perceptual and associative knowledge in category-specific impairment of semantic memory – a study of 2 cases. *Cortex*, 29: 727-740, 1993.
- LAIACONA M, BARBAROTTO R and CAPITANI E. Semantic category dissociations in naming: Is there a gender effect in Alzheimer's disease? *Neuropsychologia*, 36: 407-419, 1998.
- LAIACONA M and CAPITANI E. A case of prevailing deficit of nonliving categories or a case of prevailing sparing of living categories? *Cognitive Neuropsychology*, 18: 39-70, 2001.
- LAIACONA M, CAPITANI E and BARBAROTTO R. Semantic category dissociations: A longitudinal study of two cases. *Cortex*, 33: 441-461, 1997.
- LAWS KR. A leopard never changes its spots. *Cognitive Neuropsychology*, 15: 467-479, 1998.
- LAWS KR. Gender affects naming latencies for living and nonliving things: Implications for familiarity. *Cortex*, 35: 729-733, 1999.
- LAWS KR. Category-specific naming errors in normal subjects: The influence of evolution and experience. *Brain and Language*, 75: 123-133, 2000.
- LAWS KR. Category-specific naming and modality-specific imagery. *Brain and Cognition*, 48: 418-420, 2002(a).
- LAWS KR. Sex differences in lexical size across semantic categories. *Personality and Individual Differences*, 36: 23-32, 2003.
- LAWS KR. 'Illusions of Normality': A methodological critique of category specific-naming. *Cortex*, 41: 2005.
- LAWS KR, EVANS JJ, HODGES JR and MCCARTHY RA. Naming without knowing and appearance without associations – evidence for constructive processes in semantic memory. *Memory*, 3: 409-433, 1995.
- LAWS KR, GALE TM, LEESON VC and CRAWFORD JR. When is category specific in Alzheimer's disease? *Cortex*, 41: 2005.
- LAWS KR, LEESON VC and GALE TM. The effect of 'masking' on picture naming. *Cortex*, 38: 137-147, 2002(b).
- LAWS KR and NEVE C. A 'normal' category-specific advantage for naming living things. *Neuropsychologia*, 37: 1263-1269, 1999.
- MARQUES JF. Category-specific naming and the importance of normal performance. *Cortex*, 41: 2005.
- MARSHALL JC and GURD JM. Category-specificity: What is the question? *Cortex*, 41: 2005.
- MCMULLEN PA and FILLITER JH. Further methodological concerns in the study of category-specific identification effects. *Cortex*, 41: 2005.
- MOSS HE, TYLER LK, DURRANT-PEATFIELD M and BUNN EM. 'Two eyes of a see-through': Impaired and intact semantic knowledge in a case of selective deficit for living things. *Neurocase*, 4: 291-310, 1998.
- MOSS HE, TYLER LK and JENNINGS F. When leopards lose their spots: Knowledge of visual properties in category-specific deficits for living things. *Cognitive Neuropsychology*, 14: 901-950, 1997.
- ROSAZZA C, ZORZI M and CAPPAS SF. A reanalysis of a case of category-specific semantic impairment. *Cortex*, 41: 2005.
- SARTORI G and LOMBARDI L. Commentary on "Illusions of Normality". *Cortex*, 41: 2005.
- TURNBULL OH and LAWS KR. Loss of stored knowledge of object structure: Implications for "category-specific" deficits. *Cognitive Neuropsychology*, 17: 365-389, 2000.

Keith R. Laws, Brain and Cognition Research Group, Division of Psychology, Nottingham Trent University, Burton Street, Nottingham, NG1 4BU, UK.
e-mail: keith.laws@ntu.ac.uk