

Physical parameters of IPHAS-selected classical Be stars

I. Determination procedure and evaluation of the results[★]

L. Gkouvelis¹, J. Fabregat¹, J. Zorec^{2,3}, D. Steeghs⁴, J. E. Drew⁵, R. Raddi⁴, N. J. Wright⁵, and J. J. Drake⁶

¹ Observatorio Astronómico, Universidad de Valencia, Catedrático José Beltrán 2, 46980 Paterna, Spain
e-mail: leonardo.gkouvelis@uv.es

² Institut d'Astrophysique de Paris, CNRS (UMR 7095), 98 bis bd. Arago, 75014 Paris, France

³ UPMC, Université Paris VI, (UMR 7095), 98 bis bd. Arago, 75014 Paris, France

⁴ Department of Physics, University of Warwick, Gibbet Hill Road, Coventry CV4 7AL, UK

⁵ Centre for Astrophysics Research, STRI, University of Hertfordshire, College Lane Campus, Hatfield, AL10 9AB, UK

⁶ Smithsonian Astrophysical Observatory, 60 Garden Street, Cambridge, MA 02138, USA

Received 30 July 2015 / Accepted 11 April 2016

ABSTRACT

We present a semi-automatic procedure to obtain fundamental physical parameters and distances of classical Be (CBe) stars, based on the Barbier-Chalonge-Divan (BCD) spectrophotometric system. Our aim is to apply this procedure to a large sample of CBe stars detected by the IPHAS photometric survey, to determine their fundamental physical parameters and to explore their suitability as galactic structure tracers. In this paper we describe the methodology used and the validation of the procedure by comparing our results with those obtained from different independent astrophysical techniques for subsamples of stars in common with other studies. We also present a test case study of the galactic structure in the direction of the Perseus Galactic Arm, in order to compare our results with others recently obtained with different techniques and the same sample of stars. We did not find any significant clustering of stars at the expected positions of the Perseus and Outer Galactic Arms, in agreement with previous studies in the same area that we used for verification.

Key words. stars: early-type – stars: emission-line, Be – stars: fundamental parameters – stars: distances – Galaxy: structure – techniques: spectroscopic

1. Introduction

Classical Be (CBe) stars are main-sequence O, B and early A-type stars whose spectra show -or have shown- Balmer and other lines in emission. They are characterised by excess continuum emission in the ultraviolet, optical and infrared spectral ranges. Both the line and continuum emission arise from recombination processes in a hot, dense circumstellar decretion disk. The formation of these disks is not yet completely understood, although fast rotation, nonradial pulsations and magnetic fields are believed to be at their origin. A recent review of the nature of CBe stars and their main characteristics is presented by Rivinius et al. (2013).

In general, CBe stars have had relatively little time to move far away from their birthplaces as they are short-lived objects – particularly those of the earlier types. Being main-sequence or slightly evolved stars, they are unlikely to be embedded in their parental clouds. In addition, they are intrinsically bright, with absolute magnitudes ranging from ~ 0 to ~ -4 , enabling their detection across the whole Galactic Plane, at least in regions not affected by heavy interstellar absorption. As such, they are tracers relevant to the investigation of the spiral structure of the Galaxy.

The INT Photometric $H\alpha$ Survey of the Northern Galactic Plane (IPHAS, Drew et al. 2005; Barentsen et al. 2014) is a 1800 deg^2 imaging survey covering the entire northern Milky Way at $|b| < 5^\circ$ in the r , i and $H\alpha$ filters, using the Wide Field Camera (WFC) on the 2.5-m *Isaac Newton* Telescope (INT) in La Palma. A first list of objects displaying $H\alpha$ emission was presented by Witham et al. (2008). Most of the objects from this list have been followed up spectroscopically, at $r < 17 \text{ mag.}$, at some point between 2005 and 2012, leading to a database of low resolution spectra of 2627 objects. Preliminary analysis reveals that about 70% of this sample are CBe stars.

The general aim of this work is to obtain the fundamental physical parameters of the newly uncovered population of IPHAS CBe stars, which significantly increases the number of these objects known in the Galaxy. In addition, we aim to use this sample to contribute to the investigation of the spiral structure of the Galaxy in the northern hemisphere, using them as tracers and the standard techniques of spectroscopic parallax to measure their distances. These sources will likely obtain moderately accurate trigonometric parallaxes from *Gaia* in the near future. Therefore, their proper characterisation would also aid their future exploitation for more detailed studies of Galactic structure with *Gaia* data.

To deal with such a large set of spectra we have developed a semi-automatic procedure to obtain the relevant physical parameters of the CBe stars, including the spectral type

[★] Tables 5 and 6 are only available at the CDS via anonymous ftp to cdsarc.u-strasbg.fr (130.79.128.5) or via <http://cdsarc.u-strasbg.fr/viz-bin/qcat?J/A+A/591/A140>




Fig. 1. **a)** Graphical explanation of the BCD (D , λ_1 , ϕ_b , ϕ_{uv}) parameters in the spectrum of the B9.5V star HR4468; **b)** determination of the BCD parameters of the emission line star IPHAS J002926.93+630450.2, with B9Ve spectral type. $\Phi_{\text{Balmer lines}}$ is the line connecting the bottom of the higher Balmer lines, and δD is the difference between the values of D obtained from $\Phi_{\text{Balmer lines}}$ and Φ_{uv} .

and luminosity class, effective temperature, interstellar extinction and absolute magnitude. We have used the techniques and calibrations of the Barbier-Chalonge-Divan (BCD) spectrophotometric system (Barbier & Chalonge 1941; Chalonge & Divan 1952).

The aim of this paper is to describe the procedure we have developed to obtain CBe star astrophysical parameters from low resolution spectra, and to validate its results by comparing them with results obtained from different standard astrophysical techniques for samples of stars they have in common. A subsequent study, in preparation, will present the determination of the astrophysical parameters for the whole sample of the IPHAS CBe stars with available spectroscopy.

The paper is structured as follows: in Sect. 2 we describe the techniques of the BCD spectrophotometric system and their implementation in a semi-automatic procedure which allows us to analyse a large number of spectra efficiently, with little human intervention; in Sect. 3 we describe the spectroscopic data used for this work; in Sect. 4 we present the work we have undertaken to validate our procedure, by comparing our results with results for two samples of stars obtained with different observational data and standard astrophysical techniques; in Sect. 5 we revisit the inferred spatial distribution of the CBe stars in the direction of the Perseus Arm ($-1^\circ < b < 4^\circ$, $120^\circ < l < 140^\circ$), and compare the results with recent studies in the literature.

2. The BCD spectrophotometric system

The BCD spectrophotometric system was developed by Barbier & Chalonge (1941), and later by Chalonge & Divan (1952). The system is based upon measurable parameters around the Balmer discontinuity (BD), in the 3200–4600 \AA spectral range. The basic parameters that describe the energy distribution around the BD are: D , the Balmer jump depth, given in dex, which is an effective temperature indicator; λ_1 , the position of the Balmer discontinuity, given in \AA as the difference from 3700 \AA , which is sensitive to the surface gravity; Φ_b , a colour gradient which represent the slope of the Paschen continuum near the BD; and Φ_{uv} , the slope of the Balmer continuum.

The BD is an easily visible spectral feature for stars ranging from early O to late F spectral types. A modern description of the BCD system, together with a presentation of its advantages compared to other spectroscopic systems, is given by Zorec et al. (2009).

In Fig. 1 we present a graphical description of how the BCD parameters are measured. The value of D is calculated at $\lambda = 3700 \text{\AA}$, as $D = \log_{10}(F_{3700+}/F_{3700-})$, where F_{3700+} is the flux of the extrapolated Paschen continuum and F_{3700-} the corresponding flux on the Balmer continuum.

To measure the λ_1 parameter, the mean spectral position of the BD is obtained as the intersection between the spectral continuum and the average of the extrapolated Paschen and Balmer continua at 3700 \AA . The line that represents the average of the Φ_b and Φ_{uv} fluxes is determined by the points $\log F_\lambda - D/2$, in the Paschen continuum region at $\lambda = 4000, 4150$ and 4300\AA and $\log F_{3700} + D/2$ in the Balmer continuum at 3700 \AA .

The parameters D and λ_1 are independent of interstellar extinction, and allow the determination of the effective temperature and surface gravity by means of the calibration given by Zorec et al. (2009). The third parameter Φ_b , which measures the slope of the Paschen continuum, is a function of both the effective temperature and the interstellar extinction, and can be used to measure the interstellar reddening by means of the expression:

$$A_V = 3.1E(B - V) = 2.1(\Phi_b - \Phi_b^0). \quad (1)$$

From Aidelman et al. (2012), where Φ_b^0 is the intrinsic gradient obtained from D and λ_1 by means of the calibration given in Chalonge & Divan (1973).

2.1. The determination of the photospheric Balmer discontinuity of Be stars

The procedure described above to determine the Balmer discontinuity depth is, however, in general not applicable to CBe stars. Due to the lower pressure of the circumstellar disc, a second Balmer jump exists for these objects in the ultraviolet region of the spectrum, which is either in emission or absorption




Fig. 2. Comparison of the Balmer jump depth calculated for 23 absorption-line B-type stars, using both the ultraviolet continuum and the Balmer lines limit as explained in the text. At the left upper part it is shown the typical error in the determination of D , 0.015 dex.

(Kaiser 1989). In the case of emission, we extract the flux at F_{3700-} as the point where the bottom of the higher order Balmer line series merges into the continuum.

In Fig. 1b we illustrate this last procedure. To obtain the flux at F_{3700-} , instead of using the extrapolated Balmer continuum Φ_{uv} , we extrapolate the line traced through the bottom of the higher Balmer lines, which is labeled $\Phi_{\text{Balmer lines}}$. In the case of non emission-line stars, both methods lead to the same D values, as shown in Figs. 1a and 2, and in Cols. 6 and 7 of Table 2. The mean of the residuals between the two different measurement procedures is 0.010 ± 0.007 dex, significantly smaller than the characteristic error of 0.015 dex associated with the measure of the D parameter in the BCD system.

In the case of CBe stars, the value of D obtained in this way is the true photospheric value, not affected by circumstellar emission or absorption in the Balmer and Paschen continua. The validity of this statement is fundamental for the applicability of the BCD system to the determination of the physical parameters of CBe stars, and hence in Appendix A we discuss this issue in detail.

2.2. Setting the spectral resolution

The original calibration of the stellar parameters in the BCD system was made empirically from spectra with a mean resolution of $\Delta\lambda \sim 8 \text{ \AA}$ at the BD. Although the D parameter is independent of the resolution, the mean position of the BD, and hence the λ_1 parameter, varies with the spectral resolution. This effect is illustrated in Fig. 3, where we present the spectrum of the B-type star HR4486 at different resolutions.

Because the determination of astrophysical fundamental parameters proceeds with calibrations obtained with the original BCD parameters, to measure λ_1 we have to reduce the resolution of our spectra to the characteristic resolution of the original BCD spectrophotometric system. This is done by convolving the spectra with a Gaussian filter of the appropriate width, so that the resolution to apply the BCD formalism is obtained by

$$R_F = \sqrt{R_I^2 + G^2} \quad (2)$$




Fig. 3. The region around the BD area of the standard B-type star HR4468. The black line represents the original spectrum, and red and green lines the spectrum convolved with Gaussian filters of 2 Å and 4 Å width respectively.

where R_F is the resolution required for the BCD method, R_I is the initial resolution of the spectra, and G is the width of the Gaussian filter.

Setting of the correct spectral resolution is a complex issue, and depends on the available set of data. In Sect. 3.3. we explain how we have selected the appropriate correction of the resolution for the spectra analysed in this work.

2.3. Absolute magnitudes and distances

Absolute magnitudes in the Johnson V band (M_V) are directly obtained from the D and λ_1 parameters by means of the calibration given in Table 3 of Zorec & Briot (1991). They are converted to absolute IPHAS r magnitudes using the intrinsic ($V - R_C$) colours for dwarfs and giants supplied by Fabregat (in preparation), and assuming that Cousins R_C and IPHAS r magnitudes in the Vega system are identical within the errors involved in our procedure.

Distances are obtained by means of the standard spectroscopic parallax techniques, by comparing the absolute M_r magnitudes with the observed r magnitudes supplied in the IPHAS Second Data Release (IPHAS DR2, Barentsen et al. 2014), corrected for the interstellar absorption and the circumstellar excess due to the added flux of the disk emission.

Intrinsic r magnitudes corrected for interstellar absorption were computed using the relationship $A_r = 0.84 A_V$, from Fiorucci & Munari (2003). To correct for CBe circumstellar continuum emission we followed the method described in Sect. 3.3 of Raddi et al. (2013), which follows the earlier work by Dachs et al. (1988) that investigated the correlation between $EW(H\alpha)$ and the circumstellar colour excess $E^{cs}(B - V)$. The relations adopted in this work are

$$E^{cs}(B - V) \approx 0.02 \frac{EW(H\alpha)}{-10 \text{ \AA}}, \quad (3)$$

and

$$f_D = \frac{F_D}{F_D + F^*} \approx 0.1 \frac{EW(H\alpha)}{-30 \text{ \AA}}. \quad (4)$$

From these values we obtain the circumstellar emission in the r band, Δr , by interpolating from Table 5 of Raddi et al. (2013). The Δr correction applied for each source is given in Table 6.

2.4. Code principles

In order to analyse a large sample of spectra in a reasonable amount of time we have developed a semi-automatic procedure, only requiring user interaction to evaluate output diagrams at some important steps during the treatment of each spectrum. The basic inputs of the programme are the observed spectra, with the wavelength scale in Å and the flux in $\text{erg s}^{-1} \text{cm}^{-2}$, and a list with the names of the files containing each spectrum.

The pipeline used to treat each spectrum requires a series of inputs that the user provides as the analysis proceeds. The first step of the procedure is to ask the user for the dispersion of the spectra, and for the width G , in Å, of the Gaussian filter that the spectra require in order to transform their resolution to the original resolution of the BCD system. This first step is valid for the whole run and applies to all the spectra in the list. The next step is the analysis of the spectra one by one. As each spectrum is presented for analysis, it is first thoroughly checked to establish its wavelength range and to detect possible recording flaws.

One of the steps which requires user interaction is the wavelength scale, whose possible variations, caused either by errors in the wavelength calibration or by the radial velocity of the star, may affect the position of the Balmer discontinuity and the determination of BCD parameters. The largest shifts in the wavelength calibration were found to range from 3 to 10 Å.

A flux normalization is applied at $\lambda 4600$ Å, and the spectral coordinates transformed to $1/\lambda - \log(F/F_{4600})$, to measure the gradient Φ_b consistently with the definition of colour gradient (Allen 1973). This also enables a first estimate of the BD depth, D_1 (dex), which is considered to be preliminary because the Paschen continuum cannot be well represented by a straight line in these coordinates. D_1 is used to derive an approximate value of the stellar effective temperature. The spectrum is subsequently divided by a Planck function at this temperature. The Paschen continuum in the normalised, divided spectrum closely approach to a straight line, and can be extrapolated without ambiguity. The determination of D as the ratio of fluxes at 3700 Å removes the contribution of the Plank function.

To measure the mean spectral position of the BD we calculate the line determined by the points $\log F_\lambda - D/2$ at $\lambda = 3700, 4000, 4150$ and 4300 Å, where F_λ is the flux of the extrapolated Paschen continuum. We characterise the spectral continuum as a parabolic fit to the pseudo-continuum in the 3700–3850 Å region. The intersection between these two lines determines the value of the λ_1 parameter.

For each spectrum, a series of diagrams are constructed, with which the user can interact to control the progress of the analysis. The steps in the algorithm are shown in Fig. 4. The output of the program is a log file containing the parameters calculated for each spectrum: name, BCD(D, λ_1), T_{eff} , spectral type and luminosity class, Φ_b , Φ_b^0 , $E(B - V)$ and M_V .

3. Observations and selection of the data

3.1. IPHAS bright sample with FLWO/FAST

Follow-up spectroscopy of the emission line objects photometrically detected by IPHAS (Witham et al. 2008) was performed during the years 2005 to 2012 at the 1.5 m Fred Laurence Whipple Observatory (FLWO) Tillinghast telescope on Mount Hopkins in Arizona, using the FAST spectrograph (Fabricant et al. 1998). The data were taken with the 300 lines mm^{-1} grating, and a projected slit width of 3". The data span a wavelength range from 3500 to 7500 Å at a spectral resolution of $\Delta\lambda \approx 6$ Å.




Fig. 4. Simplified block diagram of the spectral analysis pipeline based on the BCD classification system presented in this work.

The data were processed at the Telescope Data Center at the Smithsonian Astrophysical Observatory. The spectra were delivered without flux calibration. As explained in Sect. 2, the BCD method requires at least an accurate relative flux calibration, especially in the blue and near ultraviolet part of the spectrum, around the Balmer discontinuity. This spectral region is very hard to observe in a flux-calibrated way, due to the weakness of incandescent flat field calibration lamps, CCD efficiency and optical coating properties. For this reason we took special care in performing the flux calibration.

For each different night we selected calibration spectra from the FAST archive, to ensure that all spectra were calibrated with flux standards observed the same night. The calibration has been done using standard image reduction and analysis facility (IRAF) routines.

For 31 stars we have two spectra obtained at different epochs, and for one more three spectra. In Appendix B we present overplots of the different spectra obtained for each star and a detailed discussion on the reliability of the relative flux calibrations and the spectral parameters derived from them.

Table 1. BCD standard stars observed with the INT and NOT telescopes.

Name	Telescope	Sp/LC _{MK}	Sp/LC _{BCD}	λ_1 (Å)	D (dex)	$\Delta\lambda$ (Å)	Disp.(Å/pix)	G (Å)
HR 533	INT	B2 V	B2 V	65	0.144	4	1.85	4.8
HR 1122	NOT	B5 III	B5-6 III	42	0.286	2	0.72	3.5
HR 2347	NOT	B9 V	B9 V	64	0.422	2	0.72	3.5
HR 2461	INT	B8 III	B7 III	41	0.340	3	1.40	4.0

Notes. Columns 4–6 are the standard BCD spectral type and D and λ_1 parameters respectively, from Zorec & Briot (1991). The different $\Delta\lambda$ and dispersion values for the two stars observed at the INT are due to different instrumental configurations. The last column lists the width of the Gaussian filter used to reproduce the standard BCD parameters from each spectra.

We estimate the mean error of the flux calibration as the difference between the individual values of the flux divided by the mean value. The mean error in the flux of the calibrated and normalised spectra amounts to $5.1 \pm 4.5\%$ at 4000 Å, and to $15 \pm 14\%$ at 3600 Å. The last value in the ultraviolet continuum is significantly larger, as expected. However, as explained in Sect. 2.1, to obtain the lower limit of the Balmer discontinuity of the CBe stars we don't use the extrapolated ultraviolet continuum, but instead the bottom of the higher Balmer lines. Hence, even large errors in the flux calibration short of the Balmer discontinuity, which eventually could lead to an inaccurate determination of the slope or the position of the Balmer continuum, will not have any impact in the determination of the CBe stars physical parameters used through this work.

The mean difference between the measured D and λ_1 parameters from different spectra of the same star are 0.032 dex and 7.3 Å respectively. These differences are of the same order as the ranges in D and λ_1 spanned by one spectral subtype and one luminosity class respectively, as can be seen in Fig. 10 of Zorec et al. (2009).

For this work we selected the sources with spectral features characteristic of OB-type stars. Among them we selected the spectra which have $S/N \geq 30$ around the Balmer discontinuity (~ 3700 Å), where the BCD parameters are measured. Finally, we rejected the spectra in which the continuum around the BD is not well defined. This rejection was applied automatically each time that the χ^2 of a parabolic fit to the pseudo-continuum drawn in the BD region is larger than 3σ as compared to the average of fits obtained for the entire stellar sample. We considered that in those cases we could not derive a reliable λ_1 parameter.

We examined an initial sample of 612 FAST spectra in the Perseus Arm region. After rejecting the spectra whose characteristics did not correspond to an OB-type star, and those not meeting the criteria presented in the previous paragraph, we kept a final sample of 257 spectra for further analysis.

3.2. INT and NOT spectra

Mid-resolution (2–4 Å) and high S/N (30–100 at 3700 Å) spectra of 67 CBe stars were obtained at the Roque de los Muchachos Observatory in La Palma, Canary Islands, Spain. The telescopes and instruments used were the *Isaac Newton* Telescope (INT) equipped with the Intermediate Dispersion Spectrograph (IDS), and the Nordic Optical Telescope (NOT), using the Andalucia Faint Object Spectrograph and Camera (ALFOSC). A sample of spectrophotometric standard stars, for relative flux calibration, and MK standard stars, were also observed. A complete description of this data sample is given in Sect. 2.3 of Raddi et al. (2013).

These spectra have already been analysed by Raddi et al. (2013). In this work we will use them in the procedure to determine the value of G , the width of the gaussian filter required for the FLWO/FAST spectra, as described in the next subsection, and to compare the results of our analysis with those obtained by Raddi et al. (2013) via energy distribution fitting to appropriate model atmospheres.

3.3. Gaussian filtering

For every dataset, with different resolution, we must find the appropriate gaussian width with which to convolve the spectra in order to achieve the desired BCD resolution. The best way of doing this is to observe BCD standard stars with the same instrumental configuration as the programme stars. A list of BCD standards, that is, a sample of stars for which standard values of the D and λ_1 parameters are known, is given by Zorec & Briot (1991).

No BCD standards were observed with the FAST spectrograph, and only two were obtained at the INT and two more at the NOT. In order to determine the value of G (Eq. (2)) and correct the resolution for all spectra we take the following steps:

1. As a starting point we used the INT and NOT spectra of the stars for which the standard D and λ_1 are known. For each star we computed the BCD parameters using Gaussian filters of different width, and compared the obtained values with the standard ones. The best agreement for each star was obtained with the Gaussian widths presented in Table 1.
2. We obtained the BCD parameters of all MK standard stars observed with the INT and NOT, using the widths in Table 1. From the BCD parameters we obtained the spectral types and luminosity classes, and compared them with the standard MK ones. This comparison is presented in Table 2. The agreement between the MK and BCD classification is fairly good, within two spectral subtypes and one luminosity class for most of the stars. When comparing the classification of stars in both systems, we have to keep in mind that the MK system assigns discrete spectral types, where each encompasses comparatively large intervals of (D, λ_1) parameters. Obvious differences can then appear when interpreting continuous runs of BCD parameters with discrete MK assignments.

From the above procedure, we find $G = 4.8, 4.0$ and 3.5 Å as the widths of the Gaussian filters to reduce the resolution of the INT (with two different instrumental configurations) and NOT spectra, respectively. We subsequently applied Gaussian filters of these widths to the whole sample of INT and NOT spectra, and computed the BCD parameters for all of them.

Table 2. Sample of 23 MK standard stars observed with the INT and NOT telescopes, covering almost all the B sub-types.

Name	Telescope	Sp/LC _{MK}	Sp/LC _{BCD}	λ_1 (Å)	D_{uv} (dex)	$D_{B,I}$ (dex)	$ D_{uv} - D_{B,I} $
HR 533	INT	B2 V	B2 V	79.4	0.162	0.149	0.013
HR 927	INT	B8 V	B7-8 V	54.3	0.346	0.345	0.001
HR 1122	NOT	B5 III	B5-6 III	36.1	0.236	0.237	0.001
HR 1399	INT	B5-6 V	B6 V	68.5	0.287	0.295	0.008
HR 1497	INT	B3 V	B7-8 V	77.0	0.294	0.297	0.003
HR 1576	INT	B9 V	B7 V	67.9	0.283	0.288	0.005
HR 1595	NOT	B2 V	B1 V	63.6	0.126	0.116	0.010
HR 1760	INT	A3V	>A2	74.9	0.545	0.528	0.017
HR 1808	NOT	B5 V	B4 V	62.1	0.260	0.248	0.012
HR 1820	NOT	B2 V	B3 V	60.3	0.189	0.193	0.004
HR 1860	NOT	B6 V	B6 V	60.5	0.317	0.301	0.016
HR 1863	NOT	B2.5 V	B4 V	69.0	0.222	0.210	0.012
HR 1892	NOT	B1 V	B0 V	61.4	0.104	0.105	0.001
HR 2010	NOT	B9 IV	A0-1 V	65.7	0.483	0.464	0.019
HR 2116	NOT	B8 V	B7-8 V	62.7	0.355	0.340	0.015
HR 2161	NOT	B3 V	B3 V	64.4	0.205	0.213	0.008
HR 2344	INT	B2 V	B3 IV	54.0	0.183	0.182	0.001
HR 2347	NOT	B9 V	A0 V	66.6	0.473	0.453	0.020
HR 2461	INT	B8 III	B7-8 IV	45.4	0.349	0.351	0.002
HR 2490	NOT	B3 IV	B3 IV	51.6	0.242	0.227	0.015
HR 2840	INT	B6 IV	B7 IV	47.7	0.336	0.351	0.015
HR 7996	INT	B3 III	B3-4 IV	48.9	0.270	0.252	0.018
HR 8403	INT	B5 III	B5 V	54.3	0.331	0.313	0.018

Notes. We present the measured BCD parameters and the comparison between the standard MK type and the obtained BCD spectral classification.

- The final step is to set the resolution for the FAST spectra. The procedure is described graphically in Fig. 5. We convolved the FAST spectra, one by one, starting from a Gaussian width of 1.5 Å up to 8 Å, with a 0.5 Å step. For all the stars in common between the INT/NOT and FAST samples we plot the difference $|\lambda_{1,INT/NOT} - \lambda_{1,FAST}|$ versus the Gaussian width. The minimum difference is found at $G = 4.25$ Å, and we assume this value to be the Gaussian width, G , with which to convolve all the FAST spectra prior to the determination of the BCD parameters from them.

4. Evaluation of the procedure

We now validate the described procedure, by comparing our results with results from the literature obtained for the same stars with different astrophysical techniques. For this comparison, we used the results obtained from spectroscopic data analysis by Raddi et al. (2013), and from Strömgen photometry calibration of data presented by Fabregat & Capilla (2005) and Monguió et al. (2013).

4.1. Validation against previous spectroscopic analyses

Raddi et al. (2013) studied a group of 67 candidate CBe stars in the region of the Perseus Arm, by analysing the mid resolution spectra obtained with the INT and NOT telescopes at La Palma described in Sect. 3.2. They determined their spectral types and measured their colour excess via spectral energy distribution fitting to appropriate model atmospheres in the blue part of the spectrum (3800–5000 Å).

For this comparison we use 35 spectra that meet the selection criteria described in Sect. 3.1. The results are presented




Fig. 5. Differences in the λ_1 parameter obtained with the INT/NOT spectra and the FAST spectra convolved with different Gaussian widths ranging from 1.5 Å to 8 Å, at steps of 0.5 Å, for a subset of the stars in common between the two samples. Each colour represents a spectrum and the minimum of the $\Delta\lambda_1$ difference corresponds to the convolution that best fits the INT/NOT spectrum.

in Table 3. To provide an additional element of comparison, in Table 3 we also present spectral types and luminosity classes obtained from the INT/NOT spectra by means of the standard MK classification procedures, using only the strength and widths of the spectral lines, and the ratios between lines, such as temperature and luminosity criteria, as described for example by Gray & Corbally (2009). In the last two columns we present the $E(B - V)$ obtained from the measured BCD parameters by

Table 3. BCD parameters and classification of 35 IPHAS CBe stars with INT/NOT spectroscopy, and comparison with MK classification and spectral classification given by Raddi et al. (2013).

Name	D (dex)	λ_1 (Å)	Φ_b	Φ_b^0	Sp/LC _{BCD}	Sp/LC _{MK}	Raddi et al.	$E(B - V)$	M_V
J002441.73 + 642137.5	0.291	36.5	2.85	0.80	B5-6III	B5III	B5III	1.38	-1.9
J002926.93 + 630450.2	0.355	72.3	1.51	0.85	B9V	B5-8V	B7V	0.44	1.0
J004014.89 + 651644.0	0.174	79.4	2.26	0.73	B3V	B2-3V	B2V	1.03	-1.5
J005029.25 + 653330.8	0.280	53.5	2.41	0.78	B5-6V	B3-4V	B7IV	1.10	-0.8
J005436.84 + 630549.9	0.195	63.3	2.10	0.73	B3V	B3V	B2-3V	0.92	-1.4
J005611.62 + 630350.5	0.280	69.9	1.53	0.79	B7V	B5-6V	B5V	0.50	-0.1
J005619.50 + 625824.0	0.288	70.2	1.07	0.80	B7V	B3-5V	B5V	0.18	-0.1
J010707.68 + 625117.0	0.291	60.8	2.40	0.79	B5V	B6-7Ia	B5V	1.09	-0.1
J012405.42 + 660059.9	0.246	54.8	1.94	0.76	B4V	B2-3V	B6IV	0.79	-1.0
J012751.29 + 655104.0	0.201	63.5	2.02	0.73	B3V	B3V	B7V	0.87	-1.3
J014458.14 + 633244.0	0.346	63.9	1.92	0.82	B7-8V	B3-4V	B7IV	0.74	0.6
J014620.44 + 644802.5	0.315	54.8	2.02	0.80	B6V	B5V	B7V	0.82	-0.4
J014905.18 + 624912.3	0.111	34.5	2.98	0.70	B2Ib	B2IV-III	B3IV	1.54	-5.6
J015037.67 + 644446.9	0.203	64.1	1.87	0.73	B3-4V	B3-5V	B4V	0.77	-1.1
J015246.27 + 630315.0	0.403	55.9	2.02	0.86	B9V	B8-9V	B8-9III	0.78	0.1
J015613.22 + 635623.8	0.165	68.8	1.64	0.72	B2V	B2V	B3V	0.62	-2.0
J015918.32 + 654955.8	0.321	69.0	2.22	0.85	B8V	B7-8V	B6IV	0.92	0.5
J015922.53 + 635829.3	0.234	57.1	1.91	0.75	B4V	B2-3V	B2-3V	0.78	-0.9
J022337.05 + 601602.8	0.185	52.9	2.26	0.73	B3-4IV	B3-4V	B7IV	1.03	-2.2
J023031.39 + 594127.1	0.268	50.0	1.95	0.77	B5-6V	B6-7V	B9V	0.79	-1.0
J023404.70 + 605914.4	0.184	60.5	2.71	0.73	B3-4V	B2V	B3IV	1.34	-1.5
J023642.66 + 614714.9	0.209	47.2	1.82	0.75	B3-4IV	B1-3V	B5V	0.72	-2.0
J023744.52 + 605352.8	0.142	43.6	2.52	0.70	B2III	–	B8	1.23	-3.7
J024054.96 + 630009.7	0.214	50.3	1.99	0.74	B3-4IV	B1-3V	B6	0.84	-1.6
J024146.74 + 602532.2	0.176	52.4	2.37	0.72	B2IV	B3III	B7V	1.11	-2.2
J024159.21 + 600106.0	0.197	57.6	2.18	0.73	B3-4V	B5V	B5V	0.98	-1.6
J024317.68 + 603205.5	0.120	44.2	2.26	0.70	B1III	B1.5-2III	B7V	1.05	-4.3
J024504.86 + 612502.0	0.257	43.4	2.09	0.78	B5-6IV	B6	B7IV	0.88	-1.8
J024618.12 + 613514.7	0.171	59.8	1.79	0.72	B2V	B2V	B3V	0.72	-1.9
J025016.66 + 624435.6	0.345	44.4	1.77	0.83	B7-8IV	B8-9V	B8-9III	0.63	-0.9
J025059.14 + 615648.7	0.147	62.7	1.76	0.70	B2V	B3V	B3-4V	0.71	-1.9
J025233.25 + 615902.2	0.244	40.7	1.83	0.77	B5-6III	B4V	B7V	0.71	-2.1
J025448.85 + 605832.1	0.296	59.5	1.97	0.79	B5-6V	B6-7V	B6V	0.79	-0.2
J025610.40 + 580629.6	0.161	55.2	2.04	0.71	B2V	B3-4V	B5V	0.90	-2.4
J025700.49 + 575742.8	0.152	47.1	2.24	0.71	B2III	B3-4V	B4V	1.03	-3.2

means of Eq. (1), and the absolute magnitudes in the scale defined by Zorec & Briot (1991). We estimate a mean error of 0.05 mag. for the $E(B - V)$ determination, obtained from the standard propagation of the errors through Eq. (1), considering the characteristic uncertainties of the BCD parameters involved. The mean uncertainty of the M_V values amounts to 0.15 mag. (Zorec & Briot 1991).

From Table 3 we can see a general good agreement within the three classification systems. When comparing our results with those of Raddi et al. (2013) we find that 29 of the 35 stars agree within one or two spectral subtypes. In the remaining six stars, however, there are large differences. Better agreement, with differences not larger than two sub-spectral types, is found between the BCD results and the MK classification applied to the same spectra.

In Fig. 6 we present the comparison between the $E(B - V)$ obtained by Raddi et al. (2013) and our procedure. A mean error of 0.05 mag. for our determination of the $E(B - V)$ is assumed, as stated above. For the values of Raddi et al. (2013) we used the errors quoted by these authors. There is a good agreement

between the two sets of data, with a mean difference of 0.04 ± 0.15 mag.

4.2. Validation with $uvby\beta$ photometry

Monguió et al. (2013) present a catalogue of $uvby\beta$ Strömgren-Crawford photometry for 35974 stars in the galactic anticentre direction. Thirteen classical Be stars for which we have FLWO/FAST spectroscopy have photometric data in this catalogue. In addition, two more CBe stars in the area of the galactic open cluster NGC 663, with available spectroscopy, have $uvby\beta$ photometry published by Fabregat & Capilla (2005).

For these stars we have obtained the interstellar reddening, spectral classification and absolute magnitude, using the $uvby\beta$ photometry calibrations given by Crawford (1978), Balona & Shobbrook (1984) and Moon (1986). We have followed the procedures described by Fabregat & Torrejón (1998) to apply the above calibrations to CBe stars, for which the circumstellar emission in the Balmer and Paschen continua and in the H β line have large effects on the photometric indices.

Table 4. Comparison between the spectral classification and physical parameters obtained for 15 CBe stars and those obtained from Strömgren-Crawford photometry calibrations.

Name	$D(\text{dex})$	$\lambda_1(\text{\AA})$	Sp/LC _{BCD}	Sp/LC _{<i>uwby</i>\beta}	$E(B-V)_{\text{BCD}}$	$E(B-V)_{\text{uwby}\beta}$	$M_{V,\text{BCD}}$	$M_{V,\text{uwby}\beta}$
J014602.11 + 611502.2	0.191	67.2	B4V	B5-6V	0.917	1.113	-0.3	-0.5
J014624.42 + 611037.3	0.225	68.2	B5V	B5III	0.758	0.774	-0.7	-2.3
J053237.14 + 260107.3	0.290	47.9	B5-6V	B4V	0.917	0.990	-1.4	-1.6
J053513.10 + 295912.4	0.246	31.8	B3-4III	B3III	1.168	1.074	-2.7	-2.5
J053554.13 + 295756.4	0.221	77.9	B5V	B5V	1.362	1.290	-1.4	-0.8
J053654.85 + 301757.8	0.273	66.1	B5-6V	B5V	0.870	0.953	-0.2	-0.4
J054033.87 + 274552.4	0.398	65.4	B8V	B8V	0.619	0.758	0.5	0.2
J054115.07 + 274803.2	0.139	68.4	B2V	B3V	0.997	1.000	-1.7	-0.7
J054159.24 + 274038.1	0.254	74.6	B5-6V	B5V	0.774	0.710	-0.4	-0.4
J054200.54 + 304956.6	0.355	44.4	B7IV	B7V	0.599	0.743	-0.0	-0.1
J054450.31 + 290754.5	0.319	68.7	B7V	B7V	0.568	0.510	0.2	0.0
J054603.64 + 272729.5	0.211	48.9	B3IV	B2.5V	0.973	0.963	-1.1	-1.9
J054837.64 + 281710.3	0.179	63.4	B3V	B2V	0.945	0.843	-1.4	-0.9
J054848.26 + 283547.8	0.159	60.7	B2V	B1.5-2V	1.038	0.899	-1.5	-2.2
J054937.88 + 281123.3	0.329	70.3	B7V	B8V	0.520	0.438	0.2	0.2




Fig. 6. Comparison between the colour excesses obtained from the FAST spectra with the procedure described in this work and the values given by Raddi et al. (2013).

In Table 4 we compare the results from the *uwby* β photometry analysis with those obtained with the BCD techniques described in the previous sections. The differences in the spectral types between the two methods is within ± 1 subtype, as was the case between the BCD and MK classification systems. Regarding the luminosity class, we must note that the Strömgren photometry calibrations discriminate only between classes V and III, while the BCD procedures allow a more precise separation of classes V, IV and III.

In Fig. 7 we compare the values of $E(B-V)$ obtained with the BCD and Strömgren photometry techniques. For our determination we consider an error of 0.05 mag. as in the previous subsection. For the *uwby* β calibration we use the error of 0.03 mag. quoted in Fabregat & Torrejón (1998). The agreement is very good, with a mean difference of $=0.00 \pm 0.10$ mag. In both the *uwby* β calibrations and the BCD system the intrinsic colours of the stars are determined mainly from the depth of the Balmer discontinuity, measured through the c_1 index in the *uwby* photometric system and through the D parameter in the BCD system. Both determinations are consistent and lead to similar results.




Fig. 7. Comparison between the colour excesses obtained from the FAST spectra with the procedure described in this work and those obtained with Strömgren photometry techniques.

5. Analysis of the Perseus Arm area

As a further evaluation of our procedure, in this section we present the analysis of a sample of IPHAS follow-up spectra obtained with the FLWO/FAST spectrograph. We have selected the stars located in the Perseus arm region ($-1^\circ < b < 4^\circ$, $120^\circ < l < 140^\circ$). The population of CBe stars in this area has been investigated by Raddi et al. (2013), who analysed the INT and NOT spectra described in Sect. 3.2, and by Raddi et al. (2015) also using a sample of FAST spectra. In both papers the spectral classification was performed by comparison with standard templates, complemented with the application of standard procedures of the MK classification system. Both works stress the difficulty of assigning luminosity classes, due to the fact that the profiles of the Balmer lines, which provide the main luminosity indicators in the considered spectral range, are contaminated by the circumstellar emission characteristic of CBe stars.

In the BCD system, luminosities are derived from the λ_1 parameter, which is not affected by emission. As distance determination with the spectroscopic parallax technique heavily relies on the luminosity class of the objects, it is worthwhile to




Fig. 8. Histogram distribution of the spectral types.

reproduce the above studies with reddenings and distances computed with the techniques described in this work, in order to compare the results.

Our analysis was done with 257 FAST spectra of 224 stars, that meet the criteria presented in Sect. 3.1. From each spectrum we determined the BCD (D , λ_1) parameters, the effective temperature, spectral type and luminosity class, interstellar and circumstellar colour excesses, absolute magnitude and distance. Results are presented in Tables 5 and 6. It took on average 2–3 min of work to analyse each star by means of our semi-automatic procedure.

We estimate the internal relative error in the determination of the effective temperature and distance as the difference between the individual values obtained from different spectra of the same star divided by the mean value. The mean errors measured in this way amount to 8% in T_{eff} and 24% in distance.

Eighteen stars, representing 8% of the sample, display the $H\alpha$ line in absorption. Since all the observed stars have been selected from a list of photometrically detected emission line stars (Witham et al. 2008), this figure can be considered as a measure of the false detection percentage of the photometric method. It should be noted, however, that the Be phenomenon is variable, and for some stars the lack of $H\alpha$ emission may be due to a phase transition from emission to absorption in the interval between the acquisition of the photometric and spectroscopic data. The above false detection figure must be considered as an upper limit.

Regarding the spectral classification, we found all sub-types from O8–9 up to A2. We found one Oe star, #66, with type O8–9IVe. The presence of visible HeII lines in its spectrum confirms this classification. In Fig. 8 we present the histogram of the CBe star sample distribution as a function of the spectral types.

The spatial distribution found for the CBe star sample is presented in Fig. 9, where the 224 stars with distances listed in Table 6 are plotted. For stars with more than one spectrum we used the mean value of the distance determinations. The locations of the Perseus and Outer Galactic arms are marked with dotted lines, following the range of distances given by Russeil et al. (2007). Green represents the Perseus Arm at ~ 2 – 3.5 kpc. and blue the Outer Arm at ~ 6 – 7 kpc. The distribution of the stars does not present an apparent clustering in or around these two structures. Instead, they appear scattered along the two arms and the space in between, with some

stars spread along larger distances, beyond the expected location of the Outer Arm. In Fig. 10 we present an histogram of the measured distances. These results are consistent with the findings of Raddi et al. (2013).

For ten stars we found distances in excess of 9 kpc, well beyond the expected location of the Outer Galactic Arm. Five of them have distances larger than 11 kpc and hence they are out of the scale of Fig. 9. These distances in some cases can be due to large errors in the absolute magnitude or the spectral classification of the stars. A detailed discussion on the uncertainties and possible biases in distance determination based on absolute magnitudes was given in Raddi et al. (2013). We are, however, studying in detail, for a future work a few stars in which the large distance seems to be consistently derived from high signal to noise spectra, in order to use them as tracers of the stellar population in the outskirts of the galactic plane.

6. Conclusions

This work is the first part of a larger project to study the whole population of CBe stars photometrically detected by the IPHAS survey, and use them as galactic structure tracers. The study will be based on the analysis of follow-up spectroscopy obtained at the FLWO telescope with the FAST spectrograph.

In this paper we have presented the method devised to obtain the spectral classification and the astrophysical parameters of the stars from the FLWO/FAST spectra, using the semi-automatic procedure we developed based on the BCD (Barbier-Chalonge-Divan) spectrophotometric system.

We have validated the method by comparing its results for two samples of CBe stars with independent results for the same stars obtained with different astrophysical techniques. In particular, we compared our results with those obtained with spectral template fitting and MK classification system standard techniques for one of the samples, and with Strömgren photometry standard calibrations for the other. In both cases we obtained a general good agreement in the spectral classification, within two spectral subtypes and one luminosity class for most of the stars. We have also compared our results on the interstellar extinction with those of the two samples referred to above, obtaining a good agreement in both cases.

We have also analysed a sample of CBe stars in the direction of the Perseus Arm ($-1^\circ < b < 4^\circ$, $120^\circ < l < 140^\circ$). We didn't find any significant clustering of stars at the expected distances of the Perseus and Outer Arms. Even with over three times the number of stars considered by Raddi et al. (2013) we obtain the same negative result, indicating that the errors involved in the spectroscopic parallax and absolute magnitude determination blur any hint of Galactic spiral structure, if indeed the CBe stars trace it.

Acknowledgements. This paper made use of spectroscopic data that were obtained at the FLWO-1.5 m with FAST, which is operated by Harvard-Smithsonian Centre for Astrophysics. In particular we want to thank Perry Berlind and Mike Calkins for their role in obtaining most of the FLWO-1.5 m/FAST data. The paper also makes use of data obtained as part of the INT Photometric $H\alpha$ Survey of the Northern Galactic Plane (IPHAS) carried out at the Isaac Newton Telescope (INT). The INT is operated on the island of La Palma by the Isaac Newton Group in the Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. All IPHAS data are processed by the Cambridge Astronomical Survey Unit, at the Institute of Astronomy in Cambridge. We also acknowledge the use of data obtained at the INT and the Nordic Optical Telescope as part of a CCI International Time Programme. The work of L.G. and J.F. is supported by the Spanish Ministerio de Economía y Competitividad, and FEDER, under contracts AYA2010-18352 and




Fig. 9. Distribution of the CBe stars distances as a function of the Galactic longitude. The green and blue dashed lines mark the expected positions of the Perseus and Outer Galactic arms, respectively. Five stars are at distances too long to appear in the diagram. The colour bar at the right-hand side represents the effective temperature of the stars.




Fig. 10. Histogram distribution of the distances.

References

- Aidelman, Y., Cidale, L. S., Zorec, J., & Arias, M. L. 2012, *A&A*, 544, A64
- Allen, C. W. 1973, *Astrophysical quantities* (London: Athlone Press)
- Balona, L. A., & Shobbrook, R. R. 1984, *MNRAS*, 211, 375
- Barbier, D. 1948, *Annales d'Astrophysique*, 11, 13
- Barbier, D., & Chalonge, D. 1939, *ApJ*, 90, 627
- Barbier, D., & Chalonge, D. 1941, *Annales d'Astrophysique*, 4, 30
- Barentsen, G., Farnhill, H. J., Drew, J. E., et al. 2014, *MNRAS*, 444, 3230
- Carciofi, A. C., Okazaki, A. T., Le Bouquin, J.-B., et al. 2009, *A&A*, 504, 915
- Chalonge, D., & Divan, L. 1952, *Annales d'Astrophysique*, 15, 201
- Chalonge, D., & Divan, L. 1973, *A&A*, 23, 69
- Cochetti, Y. R. 2014, Tesis de Licenciatura, Universidad Nacional de La Plata
- Cook, K. H., Alcock, C., Allsman, H. A., et al. 1995, in *IAU Colloq. 155: Astrophysical Applications of Stellar Pulsation*, eds. R. S. Stobie, & P. A. Whitelock, *ASP Conf. Ser.*, 83, 221
- Crawford, D. L. 1978, *AJ*, 83, 48
- Dachs, J., Kiehling, R., & Engels, D. 1988, *A&A*, 194, 167
- De Loore, C., Hensberge, H., Sterken, C., et al. 1979, *A&A*, 78, 287
- de Wit, W. J., Lamers, H. J. G. L. M., Marquette, J. B., & Beaulieu, J. P. 2006, *A&A*, 456, 1027
- Divan, L. 1954, *Annales d'Astrophysique*, 17, 456
- Divan, L., & Zorec, J. 1982a, in *Be Stars*, eds. M. Jасhek, & H.-G. Groth, *IAU Symp.*, 98, 61
- Divan, L., & Zorec, J. 1982b, in *Third European IUE Conf.*, eds. E. Rolfe, & A. Heck, *ESA SP*, 176, 291
- Divan, L., Zorec, J., & Andriolat, Y. 1983, *A&A*, 126, L8
- Drew, J. E., Greimel, R., Irwin, M. J., et al. 2005, *MNRAS*, 362, 753
- Fabregat, J., & Capilla, G. 2005, *MNRAS*, 358, 66
- Fabregat, J., & Torrejón, J. M. 1998, *A&A*, 332, 643
- Fabricant, D., Cheimets, P., Caldwell, N., & Geary, J. 1998, *PASP*, 110, 79
- Fiorucci, M., & Munari, U. 2003, *A&A*, 401, 781

AYA2013-48623-C2-2-P, and by the Generalitat Valenciana project of excellence PROMETEOII/2014/060. R.R. received funding from the European Research Council under the European Union's Seventh Framework Programme (FP/2007-2013)/ERC Grant Agreement No. 320964 (WDTracer). N.J.W. was supported by a Royal Astronomical Society fellowship. Finally, we are grateful to Dr. Pablo Reig for his assistance in the classification in the MK system.

- Gray, R. O., & Corbally, J., C. 2009, *Stellar Spectral Classification* (Princeton University Press)
- Hubert, A. M., & Floquet, M. 1998, *A&A*, **335**, 565
- Hubert, A. M., Floquet, M., & Zorec, J. 2000, in IAU Colloq. 175: The Be Phenomenon in Early-Type Stars, eds. M. A. Smith, H. F. Henrichs, & J. Fabregat, *ASP Conf. Ser.*, **214**, 348
- Hummer, D. G., & Mihalas, D. 1988, *ApJ*, **331**, 794
- Kaiser, D. 1989, *A&A*, **222**, 187
- Keller, S. C., Bessell, M. S., Cook, K. H., Geha, M., & Syphers, D. 2002, *AJ*, **124**, 2039
- Kurucz, R. 1993, ATLAS9 Stellar Atmosphere Programs and 2 km s⁻¹ grid, Kurucz CD-ROM No. 13 (Cambridge, Mass.: Smithsonian Astrophysical Observatory)
- Mennickent, R. E., Pietrzyński, G., Gieren, W., & Szewczyk, O. 2002, *A&A*, **393**, 887
- Monguió, M., Figueras, F., & Grosbøl, P. 2013, *A&A*, **549**, A78
- Moon, T. 1986, *Ap&SS*, **122**, 173
- Moujtahid, A. 1998, Ph.D. Thesis, Université Pierre & Marie Curie, Paris
- Moujtahid, A., Zorec, J., Hubert, A. M., Garcia, A., & Burki, G. 1998, *A&AS*, **129**, 289
- Moujtahid, A., Zorec, J., & Hubert, A. M. 1999, *A&A*, **349**, 151
- Moujtahid, A., Zorec, J., & Hubert, A. M. 2000a, in IAU Colloq. 175: The Be Phenomenon in Early-Type Stars, eds. M. A. Smith, H. F. Henrichs, & J. Fabregat, *ASP Conf. Ser.*, **214**, 506
- Moujtahid, A., Zorec, J., & Hubert, A. M. 2000b, in IAU Colloq. 175: The Be Phenomenon in Early-Type Stars, eds. M. A. Smith, H. F. Henrichs, & J. Fabregat, *ASP Conf. Ser.*, **214**, 510
- Raddi, R., Drew, J. E., Fabregat, J., et al. 2013, *MNRAS*, **430**, 2169
- Raddi, R., Drew, J. E., Steeghs, D., et al. 2015, *MNRAS*, **446**, 274
- Rivinius, T., Carciofi, A. C., & Martayan, C. 2013, *A&ARv*, **21**, 69
- Rohrmann, R. D., & Zorec, J. 2006, *Phys. Rev. E*, **74**, 041120
- Russeil, D., Adami, C., & Georgelin, Y. M. 2007, *A&A*, **470**, 161
- Vinicius, M. M. F., Zorec, J., Leister, N. V., & Levenhagen, R. S. 2006, *A&A*, **446**, 643
- Witham, A. R., Knigge, C., Drew, J. E., et al. 2008, *MNRAS*, **384**, 1277
- Zorec, J. 1986, Thèse d'État, Université Paris VII
- Zorec, J., & Briot, D. 1991, *A&A*, **245**, 150
- Zorec, J., Divan, L., & Hoefflich, P. 1989, *A&A*, **210**, 279
- Zorec, J., Arias, M. L., Cidale, L., & Ringuelet, A. E. 2007, *A&A*, **470**, 239
- Zorec, J., Cidale, L., Arias, M. L., et al. 2009, *A&A*, **501**, 297

Appendix A: The determination of the BCD parameters in CBe stars

After the discovery of the two-component Balmer discontinuity in ζ Tau by [Barbier & Chalonge \(1939\)](#), the long series of observations carried on classical Be stars in the BCD system revealed that this feature is a common phenomenon among CBe stars. The so-called first, or photospheric-like component of the BD, resembles that of a normal B-type star, whilst the second one is in emission or in absorption relative to the normal level of the Balmer continuum, depending on the Be-phase of the object. The strongest absorption or emission in this second component of the BD always produces at $\lambda \simeq 3647 \text{ \AA}$, close to the theoretical limit of the Balmer-line series, which indicates that it is formed in an environment where the gas pressure is significantly lower than in the stellar photosphere.

Except for γ Cas (O9Ve, HD 5394), during its huge emission episodes or outbursts, in 1935 and 1938 ([Barbier 1948](#)), in all other CBe stars observed more or less regularly, no changes have been detected in the value of the first component of the BD (D_*) within the limits of uncertainties that characterize the BCD system (0.015 dex in D and 1–2 \AA in λ_1). This has been proved, in particular, in some iconic CBe stars in the northern hemisphere as X Per (O9.5Ve, HD 24534), Pleione (B8IV-Ve, HD 23862), ζ Tau (B2IIIe, HD 37202), 48 Lib (B3-4IVe, HD 142983), 88 Her (B6IVe, HD 162732), χ Oph (B0Ve, HD 148184), and 59 Cyg (B1Ve, HD 200120) ([De Loore et al. 1979](#); [Divan et al. 1983](#); [Zorec 1986](#); [Zorec et al. 1989](#); [Divan & Zorec 1982b,a](#); [Zorec & Briot 1991](#)), but also in some frequently observed CBe stars in the southern hemisphere, in particular α Eri (B5III3, HD 10144) ([Vinicius et al. 2006](#); [Cochetti 2014](#)). The constancy of D_* has also been proved in stars that underwent Be \rightleftharpoons B \rightleftharpoons Be-shell-phase changes such as γ Cas, Pleione, 88 Her and 59 Cyg.

Calling total BD the quantity $D = D_* + \delta D$ (dex), the spectrophotometric variations of Be stars are currently different according to the CBe phases. Increasing emissions phases ($\delta D < 0$) are generally accompanied by a brightening and a reddening of the Paschen continuum, while shell phases ($\delta D > 0$) are generally characterized by little decreases of the stellar brightening, if any, but almost no change in the colour of the Paschen continuum. An extensive compilation of these spectrophotometric behaviours was published by [Moujtahid et al. \(1998\)](#).

As noted earlier in this appendix, the photospheric component of the BD defined in the BCD system corresponds to the ratio between an upper flux, $F_{\lambda_D}^+$, and a lower radiation flux, $F_{\lambda_D}^-$ determined at $\lambda_D \simeq 3700 \text{ \AA}$. It is worth insisting on some characteristics of these two fluxes when dealing with CBe stars:

- 1) $F_{\lambda_D}^+$ is an extrapolated flux of the Paschen continuum, whose purpose is to represent the energy distribution that would exist if the wings of the Stark-broadened Balmer lines Hn with $n > 8$ would not overlap each other to end up producing a pseudo-continuum energy distribution in low resolution spectra well before $\lambda_D = 3648 \text{ \AA}$;
- 2) The pseudo-continuum thus formed rejoins the level of the Balmer continuum just shortly after the BD in normal O-, B-, A-, F-type stars, that is, objects without extra emission or absorption produced by circumstellar gaseous environments;
- 3) $F_{\lambda_D}^-$ belongs to the aforementioned pseudo-continuum, which thus also lies on the Paschen continuum. It does mean that both fluxes $F_{\lambda_D}^\pm$ are affected by the same amount of emission or absorption, ΔF_{λ_D} , raised in the circumstellar envelope (or disc) by the electron scattering, bound-free and free-free

transitions mostly of hydrogen and helium atoms. This is the main reason that justifies our care to identify the point where the last lines of the Balmer series overlap. In this way, the circumstellar emission or absorption perturbing the Balmer side of the stellar continuum has no incidence on the determination of the photospheric BD, D_* .

The $F_{\lambda_D}^-$ can be affected by the emission in the highest members of the Balmer line series, but this phenomenon occurs rarely; it happened perhaps in γ Cas during its historic outbursts that we mentioned earlier. However, the empirical determination of D_* cannot avoid two effects related with the presence of the circumstellar environment:

- a) The genuine photospheric fluxes $F_{\lambda_D}^\pm$ transform into $F_{\lambda_D}^\pm + \Delta F_{\lambda_D}$, which implies that:

$$D_*^{\text{obs}} \simeq D_*^\circ + \left(\frac{\Delta F_{\lambda_D}}{F_{\lambda_D}^+} \right) (1 - 10^{D_*^\circ}), \quad (\text{A.1})$$

where D_*° is the actual photospheric BD, and according to the models we present below, in the hottest stars where these effects can be the highest, it is $\Delta F_{\lambda_D}/F_{\lambda_D}^+ \lesssim 0.05$.

- b) Since $F_{\lambda_D}^+$ is extrapolated, it is affected by the colour changes of the Paschen continuum due to the λ^3 -dependency of the circumstellar envelope-opacity.

[Moujtahid \(1998\)](#), [Moujtahid et al. \(1999, 2000b\)](#) and [Moujtahid et al. \(2000a\)](#) have sketched and interpreted the long-term spectrophotometric changes of CBe stars to estimate the physical parameters characterizing their circumstellar envelopes. Similar, but somewhat more detailed interpretations of these behaviours, were studied in ζ Tau by [Carciofi et al. \(2009\)](#). To have an overview of the circumstellar effects on the measured D_* as well as to quantify these effects, we have produced the models shown in Fig. A.1. These calculations are based on the circumstellar disc-model discussed in [Zorec et al. \(2007\)](#), which were adapted to produce non-LTE visible continua. We oriented the disc at $i = 70^\circ$ to easily produce Be and Be-Shell phases in the visual energy distributions by changing only the disc density near the central star, which is here parametrized with the particle number-density $n = N/10^{12}$ with $[N] = \text{cm}^{-3}$.

To simplify the calculations, we have adopted the average radial temperature distribution in the circumstellar disc derived in [Moujtahid et al. \(2000a\)](#), while its density follows the power-law deduced in [Zorec et al. \(2007\)](#) that was needed to account for the F II emission lines. For the models shown in Fig. A.1 we used stellar spectra given by [Kurucz \(1993\)](#) for: $T_{\text{eff}} = 30\,000 \text{ K}$, $\log g = 4.0$ (O9V); $T_{\text{eff}} = 22\,500 \text{ K}$, $\log g = 4.0$ (B2V); and $T_{\text{eff}} = 15\,000 \text{ K}$, $\log g = 4.0$ (B5V). In these models, the spectral region near $F_{\lambda_D}^+$ is only sketched to avoid detailed calculations of the level occupation probabilities in the upper hydrogen atomic level and the corresponding truncation of partition functions ([Hummer & Mihalas 1988](#)) that would necessarily allow for the still badly-mastered non-ideal effects ([Rohrman & Zorec 2006](#)).

The spectrophotometric behaviours shown in Fig. A.1 are an excerpt of many others that can be obtained just by changing the aspect angle i . The amount of continuum emission and absorption also depends on the size and opening angle of the lemniscat-like shaped circumstellar envelope. As much as it concerns the present models, it is rather difficult to obtain shell-phase in very hot stars. On the contrary, Be and shell phases can be obtained




Fig. A.1. Changes of the energy distribution in the spectral interval $0.325 \leq \lambda \leq 0.6 \mu\text{m}$ induced by a circumstellar disc around stars of spectral type O9V, B2V and B5V. In the figure are indicated the wavelength λ_D at which is measured D_* , and the central wavelength λ_V of the V magnitude. In the column $\log n$, n is given in units of $N/10^{12}$ with $[N] = \text{cm}^{-3}$.

Table A.1. Measurement errors induced by the circumstellar emission/absorption on the estimate of the photospheric component D_* BD and on the corresponding absolute magnitude $M_V(\lambda_1, D_*)$ determination.

$\log N/10^{12}$	-1.0	-0.8	-0.6	-0.4	-0.2	0.0	+0.2
Sp.T	$D_*^{\text{obs}} - D_*$						
O8V	0.049	0.000	-0.001	-0.002	-0.005	-0.012	-0.023
B2V	0.130	0.000	-0.001	-0.002	-0.009	-0.020	-0.038
$\log N/10^{12}$	-1.6	-1.4	-1.2	-1.0	-0.8	-0.6	-0.4
B5V	0.261	0.000	0.000	0.000	-0.001	-0.002	-0.005
	δM_V						
O8V	-0.01	-0.02	-0.02	-0.04	-0.09	-0.21	-0.41
O8III	-0.01	-0.01	-0.00	-0.03	-0.07	-0.16	-0.30
B2V	-0.01	-0.02	-0.02	-0.06	-0.14	-0.30	-0.60
B2III	-0.01	-0.02	-0.02	-0.04	-0.10	-0.23	-0.45
B5V	0.00	0.00	0.00	0.00	-0.01	-0.02	-0.06
B5III	0.00	0.00	0.00	0.00	-0.01	-0.02	-0.05

easily for B2-type objects, and late type B would rather display shell aspects. According to the effective temperature among early type CBe stars, the emission in the second BD is accompanied either by a reddening or a blueing of the Paschen continuum. In late type CBe stars, the shell-absorption seems not to be accompanied by any significant change of in colour of the Paschen continuum energy distribution. In all these cases, we can note that when the emission or the absorption in the second BD vary a lot, the photospheric energy jump at the BD remains remarkably unchanged.

In Table A.1 we present the values of D_* for the unperturbed stars, as well as the errors $\delta D_* = D_*^{\text{obs}} - D_*$ committed on the D_* -determination carried by the effects a) and b) described above. Because we are not interested in giving detailed physical interpretations of the calculated effects, the parameter $\log n$ must be regarded only as a way to produce a given amount of continuum emission or absorption. We should thus pay attention only on these amounts and conclude that the

events labeled with $\log n = 0.0$ and $+0.2$ are rather extreme and rarely observed. In this respect, we note that in the compilation made by Moujtahid et al. (1998), which actually corresponds to spotted measurements concerning otherwise long-term Be-star “bumper” activity (Cook et al. 1995; Hubert & Floquet 1998; Hubert et al. 2000; Keller et al. 2002; Mennickent et al. 2002; de Wit et al. 2006), the magnitude changes are of the order of $0.1 \lesssim \Delta V \lesssim 0.2$ mag and rarely as high as $\Delta V \lesssim 0.5$ mag.

In the lower part of Table A.1 we also give the errors δM_V that δD_* values would produce on the visual absolute magnitude $M_V(\lambda_1, D_*)$. We notice that in all cases the δM_V errors are smaller than the amplitude of $M_V(\lambda_1, D_*)$ values that may concern a single average MK spectral type (Zorec & Briot 1991). Let us finally mention that errors made on the estimate of D_* can be somewhat reduced if the colour effect acting on the extrapolated $F_{\lambda_D}^+$ flux, which resembles that of an increased interstellar extinction, is taken into account by using

$$\begin{aligned} D_* &= D_*^{\text{obs}} + \Delta D_* \\ \Delta D_* &= a \times [\Phi^{\text{obs}} - \Phi(\lambda_1, D_*^{\circ})], \end{aligned} \quad (\text{A.2})$$

where $a = 0.02$, if the Paschen colour gradient is $\Phi = \Phi_{\text{pb}}$ of the BCD system. This relation was derived by the creators of the BCD system (Divan 1954), and since then applied to every determination of the BD when the ISM extinction detected by a high gradient difference $\Delta\Phi = \Phi^{\text{obs}} - \Phi(\lambda_1, D_*)$. In extreme cases $\Delta\Phi \approx 0.5 \mu\text{m}$, the use of Eq. (A.2) can reduce the largest uncertainties δD_* given in Table A.1 to the standard BCD uncertainty $\delta D \lesssim 0.015$ dex. It is finally worth noting that the change of λ_1 due to δD_* and the variation of the slope of the Paschen continuum is extremely small and thus negligible.

Appendix B: Evaluation of the flux calibration of the FLWO/FAST spectra

The determination of reliable physical parameters of stars by means of the BCD system methods requires the spectra to have at least an accurate relative flux calibration. In this appendix we

present an evaluation of the flux calibration of the FLWO/FAST sample, by comparing different spectra of the same objects. In the discussion below we analyse only the spectral region between 3700 and 4600 Å, where the BCD parameters are measured. For some spectra drops in the signal-to-noise and large differences in the flux calibrations at shorter wavelengths are apparent. However, as explained in Sects. 2.1 and 3.1, they don't have any impact in the determination of the physical parameters of the stars.

For 31 stars we have two spectra obtained at different epochs, and for one more, star 181, we have three spectra. All these spectra are presented in Figs. B.1 to B.1. For each object we present two panels. In the upper one we overplot the two- or three- flux calibrated and normalised spectra in a logarithmic scale. In the bottom panel we represent the difference between the normalised fluxes as a function of the wavelength.

From the comparison between the spectra we can divide the object sample into three groups. Group A is composed of the stars for which the two calibrated spectra overlap, indicating that the flux calibrations applied to each spectrum are consistent. It includes 14 objects, namely stars 23, 56, 68, 110, 118, 124, 161, 170, 171, 179, 181, 210, 219 and 223. Group B is composed of the stars displaying differences between the spectra, which imply differences between the flux calibrations, but the difference is a lineal function of the wavelength, as shown in the bottom panel for each star. Group B includes 16 objects, namely stars 109, 116, 127, 130, 153, 154, 164, 169, 172, 183, 185, 188, 211, 214, 217, and 218. Finally, Group C is composed by two more objects, stars 204 and 207, which show differences in the flux calibration which are not a lineal function of the wavelength.

The differences in the flux calibration present in the stars of Group B do not have any effect on the determination of the D and λ_1 parameters. If the difference between the two spectra increases linearly with the decreasing wavelength, the variation of the Paschen continuum extrapolated at 3700 Å exactly compensates the variation of the bottom of the Balmer discontinuity at the same wavelength, and the position of the discontinuity is not affected. The mean differences between the D and λ_1 parameters for stars of Group A are 28 ± 24 dex and 7.0 ± 6.0 Å respectively, while for stars of Group B these are 31 ± 22 dex and 7.4 ± 5.4 Å respectively.

These differences do, however, affect the determination of the interstellar reddening. The reddening value is derived from the Φ_b parameter, which measures the slope of the Paschen continuum. This slope is sensitive to differences in the relative flux calibration. The mean difference in the determination of $E(B-V)$ for stars in the Group A is 0.06 ± 0.04 mag., while for stars in Group B it amounts to 0.15 ± 0.10 mag. This last figure is consistent with the standard deviation obtained when comparing our $E(B-V)$ values with values in the literature for the same stars presented in Sect. 4, and can be considered as the mean error of our $E(B-V)$ determination.

For stars in Group C the flux calibrations are not consistent. This represents up to four spectra out of 65 analysed in this appendix. In addition, it should be noted that the differences between the calibrations are small, and translate into mean differences of the parameters determined from them well within 3σ of the mean errors considered in this paper.




Fig. B.1. Stars with two or more spectra. For each object the *upper panel* represents the overplot of the spectra and the *lower panel* the difference between the two spectra.




Fig. B.1. continued.




Fig. B.1. continued.




Fig. B.1. continued.